arXiv:1410.8781v1 [math.HO] 29 Oct 2014

CONSTRUCTING THE FIXED POINT OF A NON-ISOMETRIC PLANE SIMILARITY

ZHIGANG HAN AND RONALD UMBLE

Abstract. We present an algorithm for constructing the fixed point of a general non-isometric similarity of the plane.

1. Introduction

A majority of states across the USA are in the process of implementing the *Common Core State Standards for Mathematics* [1]. An essential component of the geometry standard is the study of plane similarities. To quote the Standards, "*Given two similar two-dimensional figures, describe a sequence that exhibits the similarity between them.*" [CCSS.MATH.CONTENT.8.G.A.4] If the given figures are congruent, the similarity α relating them is a distance-preserving transformation, called an *isometry,* and by the Classification Theorem for Plane Isometries, α is a translation, a rotation, a reflection, or a glide reflection [2]. On the other hand, if the given figures are not congruent, α is a non-isometric similarity, and by the Classification Theorem for Plane Similarities, α is a stretch, a stretch rotation, or a stretch reflection [2]. Nevertheless, in either case, it is easy to determine the *similarity type* of α at a glance. For a discussion of the *Similarity Recognition Problem*, see [3].

The precise similarity relating two similar non-congruent plane figures can be difficult to determine, however. While the fact that *every non-isometric similarity has a fixed point* is an important clue (see Theorem 1 below), the proof only establishes existence, and it is not immediately clear how to go about constructing the fixed point. And this is essential – once the fixed point is determined, the additional requirements of the definition follow immediately. In this article we present a straight forward algorithm for constructing the fixed point of a non-isometric similarity.

2. Notation and Definitions

Let $r \in \mathbb{R}$ and let P and Q be arbitrarily chosen points. A (plane) **similarity of ratio** $r > 0$ is a transformation $\alpha : \mathbb{R}^2 \to \mathbb{R}^2$ such that if $P' = \alpha(P)$ and $Q' = \alpha(Q)$, then $P'Q' = rPQ$. An **isometry** is a similarity of ratio 1. One can show that similarities are bijective transformations that preserve betweeness; for a proof see [3].

Let C be a point. The **stretch about** C **of ratio** $r > 0$ is the transformation $\xi_{C,r} : \mathbb{R}^2 \to \mathbb{R}^2$ with the following properties: (1) $\xi_{C,r}(C) = C$; (2) *if* $P \neq C$, *then* $P' = \xi_{C,r}(P)$ *is the unique point on* $\overrightarrow{CP}$ *such that* $CP' = rCP$. The point C is

Date: September 16, 2014.

called the **center** of the stretch (see Figure 1), and it is easy to check that $\xi_{C,r}$ is a similarity of ratio r.



Figure 1. The stretch about C of ratio 3.

Let $\theta \in \mathbb{R}$. The **rotation about** C **through angle** θ is the transformation $\rho_{C,\theta} : \mathbb{R}^2 \to \mathbb{R}^2$ with the following properties: (1) $\rho_{C,\theta}(C) = C$; (2) *if* $P \neq C$ *and* $P' = \rho_{C,\theta}(P)$, *then* $CP' = CP$ *and the angle from* $\overrightarrow{CP}$ *to* $\overrightarrow{CP'}$ *is* θ. The point C is called the **center** of rotation (see Figure 2).



Figure 2. The rotation about C through angle θ.

The **halfturn about** C is the transformation $\varphi_C : \mathbb{R}^2 \to \mathbb{R}^2$ defined by $\varphi_C(P) = \rho_{C,180^\circ}(P)$. A **dilation** about C of ratio $r > 0$ is a transformation $\delta_{C,r} : \mathbb{R}^2 \to \mathbb{R}^2$ that is either a stretch or a stretch followed by a halfturn, i.e., $\delta_{C,r} = \xi_{C,r}$ or $\delta_{C,r} = \varphi_C \circ \xi_{C,r}$. The point C is the **center** of dilation, and a dilation of ratio 1 is either the identity ι or a halfturn. Furthermore, $\delta_{C,r}$ is a similarity of ratio r since the composition of a similarity of ratio r with a similarity of ratio 1 is a similarity of ratio r. When C is **between** points P and Q we write $P - C - Q$. Note that if $C - P - Q$, there is a real number $r > 0$ such that $Q = \xi_{C,r}(P)$, and if $P - C - Q$, there is a real number $r > 0$ such that $Q = \left(\varphi_C \circ \xi_{C,r}\right)(P)$. Thus in either case, $Q = \delta_{C,r}(P)$.

Let m be a line. The **reflection in line** m is the transformation $\sigma_m : \mathbb{R}^2 \to \mathbb{R}^2$ with the following properties: (1) *If* P *is on* m, *then* $\sigma_m(P) = P$; *(2) if* P *is off* m *and* $P' = \sigma_m(P)$, *then* m *is the perpendicular bisector of* $\overline{PP'}$. The line m is the **axis** of reflection (see Figure 3). A **stretch rotation** is a non-trivial stretch about a point C followed by a non-trivial rotation about C. A **stretch reflection** is a non-trivial stretch about a point C followed by a reflection in some line through C.



Figure 3. Line m is the axis of reflection.

Lines l and m are **parallel** if $l = m$ or $l \cap m = \varnothing$. A **collineation** is a transformation $\phi : \mathbb{R}^2 \to \mathbb{R}^2$ that sends lines to lines; a collineation is a **dilatation** if it sends each line to a line parallel to it. In fact, similarities are collineations and every dilatation that is a non-isometric similarity is a dilation.

3. Constructing Fixed Points

The fixed point of a non-trivial dilation $\delta_{C,r}$, namely its center C, is easy to construct. Choose any point A. If $A = \varphi_C(A)$, then $A = C$. So assume that $A \neq A' = \varphi_C(A)$. If $r = 1$, then $\delta_{C,r}(A) = \varphi_C$ and C is the midpoint of $\overline{AA'}$. If $r \neq 1$, choose any point B off $\overleftrightarrow{AA'}$. If $B = \delta_{C,r}(B)$, then $C = B$. So assume that $B \neq B' = \delta_{C,r}(B)$. Since dilations are dilatations, $\overleftrightarrow{AB} \parallel \overleftrightarrow{A'B'}$. On the other hand, $\overleftrightarrow{AA'} \nparallel \overleftrightarrow{BB'}$; otherwise, $\square AA'B'B$ is a parallelogram and $A'B' = AB$ implies $\delta_{C,r}$ is an isometry, which is a contradiction. Furthermore, $\delta_{C,r}(\overleftrightarrow{AA'}) = \overleftrightarrow{AA'}$ since $A' \in \overleftrightarrow{AA'} \cap \delta_{C,r}(\overleftrightarrow{AA'})$, and similarly $\delta_{C,r}(\overleftrightarrow{BB'}) = \overleftrightarrow{BB'}$. Therefore $\delta_{C,r}(\overleftrightarrow{AA'} \cap \overleftrightarrow{BB'}) = \overleftrightarrow{AA'} \cap \overleftrightarrow{BB'}$ and $\overleftrightarrow{AA'} \cap \overleftrightarrow{BB'} = C$ (see Figure 4).



Figure 4. Constructing the fixed point C of a non-isometric dilation.

The proof of our general fixed point construction algorithm rests heavily on the following standard existence theorem, which we include here for completeness:

Theorem 1. *Every non-isometric similarity has a fixed point.*

Proof. Let α be a non-isometric similarity. If α is dilatation, it is a dilation and has a fixed point, namely its center constructed above. So assume that α is not a dilatation. Let l be a line that cuts its image $l' = \alpha(l)$ at the point $A = l \cap l'$ and let $A' = \alpha(A)$. If $A' = A$, then α has a fixed point and we're done. So assume $A' \neq A$. Then A' is on l' and off l. Let m be the line through A' parallel to l. Since A' is off l, the parallels l and m are distinct. Let $m' = \alpha(m)$. We claim that lines l' and

m' are distinct parallels. If not, either $l' \nparallel m'$ or $l' = m'$, and in either case there exists a point $Q' \in l' \cap m' = \alpha(l) \cap \alpha(m)$ and points Q_1 on l and Q_2 on m such that $Q' = \alpha(Q_1) = \alpha(Q_2)$. But l and m are distinct parallels, hence $Q_1 \neq Q_2$ and α is not injective, which is a contradiction. Therefore l' and m' are distinct parallels, as claimed. Let $B = m \cap m'$ and let $B' = \alpha(B)$. If $B' = B$, then α has a fixed point and we're done. So assume $B' \neq B$. Then B' is on m' and $B' \neq A'$ since A' is on l'. Note that $\overleftrightarrow{AA'} = l' \neq m' = \overleftrightarrow{BB'}$ and $\overleftrightarrow{AA'} \| \overleftrightarrow{BB'}$ by the claim. Since α is not an isometry, $\square ABB'A'$ is not a parallelogram and $\overleftrightarrow{AB} \nparallel \overleftrightarrow{A'B'}$. Let $P = \overleftrightarrow{AB} \cap \overleftrightarrow{A'B'}$ and let $P' = \alpha(P)$. If $P' = P$, the proof is complete. We consider three cases:

Case 1. $A - P - B$. Then $A' - P - B'$ since $\overleftrightarrow{AA'} \| \overleftrightarrow{BB'}$ (see Figure 5) and $A' - P' - B'$ since α preserves betweenness. Thus A', P, P', and B' are collinear. Since $\overleftrightarrow{AA'} \| \overleftrightarrow{BB'}$, we have $\angle PAA' \cong \angle PBB'$ and $\angle PA'A \cong \angle PB'B$ by the Alternate Interior Angles Theorem. Hence $\triangle APA' \sim \triangle BPB'$ (AA) so that $AP/BP = A'P/B'P$ by the Similar Triangles Theorem. Let r be the ratio of the similarity α; then $A'P' = rAP$ and $B'P' = rBP$ so that

$$\frac{A'P'}{B'P'} = \frac{rAP}{rBP} = \frac{AP}{BP} = \frac{A'P}{B'P}.$$

Since $A'P' = A'B' - B'P'$ and $A'P = A'B' - B'P$, direct substitution gives

$$\frac{A'B' - B'P'}{B'P'} = \frac{A'B' - B'P}{B'P}.$$

Then $B'P' = B'P$ by algebra, and $P = P'$ as claimed.



Figure 5. The points A', P, and B' are collinear.

Cases 2 and 3. The proofs assuming $A - B - P$ and $P - A - B$ are similar and left to the reader. □

Theorem 1 asserts that every non-isometric similarity α has a fixed point, so let's construct it. By the Classification Theorem for Plane Similarities, α is a stretch, a stretch rotation, or a stretch reflection [2], so the fixed point of α is always the center of a stretch. The fixed point of a dilation was constructed above, so assume that α is not a dilation.

We need the following technical fact:

Proposition 1. *Let α be a non-isometric similarity that is not a dilation. There exist non-degenerate triangles $\triangle PQR$ and $\triangle P'Q'R'$ such that $P' = \alpha(P)$, $Q' = \alpha(Q)$, $R' = \alpha(R)$, $\overleftrightarrow{PQ} \nparallel \overleftrightarrow{P'Q'}$, and $\overleftrightarrow{QR} \nparallel \overleftrightarrow{Q'R'}$.*

Proof. Since α is not a dilation, it is not a dilatation. Hence there exists a line q such that $q' = \alpha(q) \nparallel q$, and it follows that $q'' = \alpha(q') \nparallel \alpha(q) = q'$. Let $Q = q \cap q'$. Choose a point $P \neq Q$ on q and a point $R \neq Q$ on q'. Then P, Q, and R are distinct and non-collinear, and so are $Q' = \alpha(Q) = q' \cap q''$, $P' = \alpha(P) \in q'$, and $R' = \alpha(R) \in q''$. Hence $\overleftrightarrow{PQ} = q \nparallel q' = \overleftrightarrow{P'Q'}$ and $\overleftrightarrow{QR} = q' \nparallel q'' = \overleftrightarrow{Q'R'}$. □

Our fixed point construction algorithm now follows.

Algorithm 1. *Let α be a non-isometric similarity that is not a dilation. To construct the fixed point of α, choose triangles $\triangle PQR$ and $\triangle P'Q'R'$ that satisfy the conditions of Proposition 1 and proceed as follows:*

1. *Construct line n through R parallel to $m = \overleftrightarrow{PQ}$.*
2. *Construct line n' through R' parallel to $m' = \overleftrightarrow{P'Q'}$.*
3. *Let $D = m \cap m'$ and $E = n \cap n'$.*
4. *Construct line $a = \overleftrightarrow{DE}$* (see Figure 6).
5. *Interchange P and R and repeat steps 1–4 to construct line b.*
6. *Then $C = a \cap b$ is the fixed point of α.*



Figure 6. Constructing the fixed point C of a stretch rotation.

The proof of Algorithm 1 is a consequence of the following lemma:

Lemma 1. *Let α be a non-isometric similarity and let C be the fixed point of α given by Theorem 1. Let l and m be distinct parallel lines off C, let $l' = \alpha(l)$, and let $m' = \alpha(m)$. If $l \cap l' = D$ and $m \cap m' = E$, then C, D, and E are collinear.*

Proof. Let α be a non-isometric similarity and let C be the fixed point of α given by Theorem 1. Then α is not a dilation since $l' = \alpha(l) \nparallel l$ by assumption. Hence by the Classification Theorem for Similarities, $\alpha = \rho_{C,\Theta} \circ \xi_{C,r}$ or $\alpha = \sigma_n \circ \xi_{C,r}$,

where in the latter case, C is on n. Since l and m are distinct parallel lines off C, there exists a dilation $\delta_{C,s}$ such that $\delta_{C,s}(l) = m$. By definition, either $\delta_{C,s} = \xi_{C,s}$ or $\delta_{C,s} = \varphi_C \circ \xi_{C,s}$. The reader can easily check that $\xi_{C,s}$ commutes with $\rho_{C,\Theta}$, σ_n, and $\xi_{C,r}$. Hence $\xi_{C,s}$ commutes with α. Furthermore, φ_C commutes with $\rho_{C,\Theta}$, σ_n, and $\xi_{C,r}$. Hence φ_C commutes with α. Consequently, $\delta_{C,s}$ commutes with α and $\delta_{C,s}(l') = \delta_{C,s}(\alpha(l)) = \alpha(\delta_{C,s}(l)) = \alpha(m) = m'$. Therefore $\delta_{C,s}(D) = \delta_{C,s}(l \cap l') = m \cap m' = E$, and it follows that C, D, and E are collinear. □

Theorem 2. *Let α be a non-isometric similarity and let C be the point given by Algorithm 1. Then C is the fixed point of α.*

Proof. Referring to the notation in Algorithm 1, note that $\alpha(m) = m'$ since $\alpha(P) = P'$ and $\alpha(Q) = Q'$. Thus $\alpha(n) = n'$ since $\alpha(R) = R'$ and α maps parallel lines to parallel lines. By Lemma 1, C, D, and E are collinear. A similar argument shows that C, F, and G are also collinear. Therefore $C = \overleftrightarrow{DE} \cap \overleftrightarrow{FG}$. □

Example 1. The two triangles in Figure 7 below are related by a stretch reflection α whose fixed point C was given by Algorithm 1. To construct the axis of reflection, let P be a vertex and let $P' = \alpha(P)$. Then the axis of reflection is the bisector of $\angle PCP'$.



Figure 7. Constructing the fixed point C and axis of a stretch reflection.

4. Conclusion

Given a non-isometric plane similarity α, we have presented an algorithm for constructing the fixed point of α. Our algorithm can be easily applied using a reflecting instrument such as MIRA. Since such instruments are readily available in a typical high school classroom, our algorithm gives students of plane geometry

direct hands-on access to the fixed point of a non-isometric similarity. Hands-on activities such as this create the potential for a deeper and more complete understanding of plane similarities and their properties.

References

[1] Common Core State Standards Initiative [CCSSI], *Common Core State Standards for Mathematics*, National Governors Association Center for Best Practices and the Council of Chief State School Officers, Washington, DC, 2010.

[2] Martin, George E., *Transformation Geometry: An Introduction to Symmetry,* Springer-Verlag, New York, 1982.

[3] Umble, R. and Han, Z. *Transformational Plane Geometry,* Taylor and Francis Group, Boca Raton, FL, 2014.

Department of Mathematics, Millersville University of Pennsylvania, Millersville, PA. 17551

E-mail address: zhigang.han@millersville.edu

Department of Mathematics, Millersville University of Pennsylvania, Millersville, PA. 17551

E-mail address: ron.umble@millersville.edu